



King's Research Portal

DOI:

[10.1080/00140139.2018.1489982](https://doi.org/10.1080/00140139.2018.1489982)

Document Version

Peer reviewed version

[Link to publication record in King's Research Portal](#)

Citation for published version (APA):

Haslam, C., Kazi, A., Duncan, M., Clemes, S., & Twumasi, R. (2018). Walking Works Wonders: a tailored workplace intervention evaluated over 24 months. *Ergonomics*, 62(1), 31-41.

<https://doi.org/10.1080/00140139.2018.1489982>

Citing this paper

Please note that where the full-text provided on King's Research Portal is the Author Accepted Manuscript or Post-Print version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version for pagination, volume/issue, and date of publication details. And where the final published version is provided on the Research Portal, if citing you are again advised to check the publisher's website for any subsequent corrections.


General rights

Copyright and moral rights for the publications made accessible in the Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognize and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Research Portal

Take down policy

If you believe that this document breaches copyright please contact librarypure@kcl.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.



Walking Works Wonders: a tailored workplace intervention evaluated over 24 months

Cheryl Haslam, Aadil Kazi, Myanna Duncan, Stacy Clemes & Ricardo Twumasi

To cite this article: Cheryl Haslam, Aadil Kazi, Myanna Duncan, Stacy Clemes & Ricardo Twumasi (2018): Walking Works Wonders: a tailored workplace intervention evaluated over 24 months, Ergonomics, DOI: [10.1080/00140139.2018.1489982](https://doi.org/10.1080/00140139.2018.1489982)

To link to this article: <https://doi.org/10.1080/00140139.2018.1489982>



Accepted author version posted online: 22 Jun 2018.



Submit your article to this journal [↗](#)



Article views: 29



View Crossmark data [↗](#)

Walking Works Wonders: a tailored workplace intervention evaluated over 24 months

Author names and affiliations:

Cheryl Haslam^{a*}, Aadil Kazi^b, Myanna Duncan^c, Stacy Clemes^a, Ricardo Twumasi^d

^a School of Sport, Exercise and Health Sciences, Loughborough University, Leicestershire LE11 3TU, UK

^b NIHR Imperial Biomedical Research Centre (BRC), Faculty of Medicine, Imperial College London & Imperial College Healthcare NHS Trust, St Mary's Hospital, London, W2 1NY, UK

^c Department of Psychology, Institute of Psychiatry, Psychology and Neuroscience, King's College London, Guy's Campus, London, SE1 1UL, UK

^d Alliance Manchester Business School, The University of Manchester, D3 AMBS East, Booth Street East, Manchester, M13 9SS, UK

*Corresponding author. Telephone: +44 1509 223086

E-mail address: C.O.Haslam@lboro.ac.uk

Acknowledgements

This study was part of a 5 year research project funded by the New Dynamics of Ageing Programme (RES-353-25- 0006; RCUK, led by ESRC).

Word count: 5840 (excluding references)

Abstract

This article presents longitudinal data from 1120 participants across 10 worksites enrolled in Walking Works Wonders, a tailored intervention designed to increase physical activity and reduce sedentary behaviour. The intervention was evaluated over 2 years, using a quasi-experimental design comprising 3 conditions: tailored information; standard information and control. This study explored the impact of the intervention on objective measures (BMI, %Fat, waist circumference, blood pressure and heart rate) and self-reported measures of physical activity, sedentary behaviour, physical and psychological health. Interventions tailored to employees' stage of change significantly reduced BMI and waist circumference compared to standard and control conditions. Employees who received either a standard or tailored intervention demonstrated significantly higher work ability, organizational commitment, job motivation, job satisfaction, and a reduction in intention to quit the organization. The results suggest that adopting a tailored approach to interventions is particularly effective in terms of improving health in the workplace.

Practitioner Summary

This study describes Walking Works Wonders, a tailored intervention, which aims to encourage physical activity in the workplace. The study evaluated Walking Works Wonders over a 2 year period and demonstrated that interventions are more effective in improving health outcomes where the information is tailored to employees' stage of change.

Keywords:

Stage of change, tailored workplace intervention; physical activity, sedentary behaviour.

Introduction

We are experiencing a sitting epidemic in the workplace, with ever growing numbers of people employed in sedentary occupations spending much of their working day sitting. This is a major public health issue as sedentary behaviour is associated with an increased risk of premature mortality (Biswas et al., 2015; Wilmot et al., 2012). Research has demonstrated that, even in physically active individuals, prolonged sitting is associated with an increased risk of premature mortality (Katzmarzyk et al. 2009). Sedentary behaviour is also an established risk factor for a wide range of chronic health conditions including obesity, diabetes, cardiovascular disease and cancer (Hamilton et al. 2007; Gierach et al. 2009; Katzmarzyk et al. 2009; Lynch, 2010; van Uffelen et al. 2010; Proper et al. 2011; Wilmot et al. 2011; Chau et al. 2013).

As work provides a major contribution to sedentary time on work days, the workplace has been highlighted as a key setting for interventions designed to reduce sedentary behaviour (Kazi, 2013; Kazi et al. 2014; Mansoubi et al. 2014). Walking Works Wonders is a new tailored intervention designed to increase physical activity and reduce sedentary behaviour at work. It was evaluated over a 2 year period in 10 worksites across the UK. The longitudinal study investigated changes in physiological and psychological outcomes (measured at 6 monthly intervals) and this article presents the data across the 5 time-points: baseline, 6 months, 12 months, 18 months and 24 months.

The Walking Works Wonders intervention involves tailoring health information according to employee's readiness to change. This approach is based on the Stage of Change Model (Prochaska & DiClemente, 1982; 1983) which was originally developed within the context of smoking cessation. The model assumes that behaviour change involves movement through stages:

- i) precontemplation (resistance to recognising or modifying problem behaviour)
- ii) contemplation (thinking about changing, but not ready to act)
- iii) preparation (intending to change in the next 30 days, and/or having made plans to do so)
- iv) action (changed behaviour, no longer than 6 months ago)
- v) maintenance (changed over 6 months ago, working to consolidate gains made and avoid relapse)

According to the model, stage determines receptiveness to (and the effectiveness of) health education. Individuals in the precontemplation stage require information about the health risks associated with their current behaviour whilst those in later stages (contemplation/preparation) need practical advice on how to change their behaviour. The model has been applied to a wide range of health related behaviours including: smoking cessation (Andersen & Keller, 2002; Prochaska et al., 1993); maternal smoking cessation (Haslam, 2000; Haslam and Draper, 2000; Haslam and Lawrence, 2004); exercise (Marshall & Biddle, 2001; Kirk et al. 2003) and dietary behaviours (Povey et al. 1999).

Following the successful application of the stage of change model to community health interventions, calls were made to apply the model to workplace interventions (Haslam and Haslam, 2001; Haslam, 2002). Whysall, Haslam and Haslam (2005) developed a tool for assessing employee stage of change and used this to determine if tailored ergonomic interventions are more effective in reducing musculoskeletal disorders than standard ergonomic approaches. They demonstrated that tailored interventions were significantly more effective in changing behaviour and reducing musculoskeletal disorders in a diverse range of workplace settings (Whysall, Haslam and Haslam, 2006; 2007). Rothmore et al. (2017) investigated compensable injuries among workers who had received ergonomics advice tailored to stage of change compared to standard ergonomics advice. They

demonstrated that those given tailored advice were less likely to report a compensable injury than those given standard advice.

Walking Works Wonders was developed through extensive user engagement, with employees, managers and representatives from occupational health, and the intervention adopted a stage of change approach (Kazi, 2013). An initial evaluation of Walking Works Wonders (Kazi, 2013) examined the impact of the intervention over 12 months. This paper provides an evaluation of Walking Works Wonders over the intervention period (12 months) and a further 12 months follow-up.

Health information was provided to employees through leaflets which offered standard or tailored advice. The leaflets were supplemented by posters, physical activity challenges and interactive environmental prompts. All additional site posters/challenges/environmental prompts were standardised. Only the leaflets given specifically to each employee were tailored to stage of change. In order to test the effects of tailoring information, a quasi-experimental design was used with 3 conditions: standard information; staged information and control group. The research objectives were to:

- 1) Implement an intervention aimed at increasing physical activity and reducing sedentary behaviour in the workplace
- 2) Collect longitudinal repeated measures data (the same pre-determined outcome measures collected at baseline) at 6 months (mid-intervention) 12 months (end of the intervention) plus follow-up measures at 18 and 24 months to evaluate the long term effectiveness of the intervention
- 3) Identify whether significant differences exist between changes in the outcome measures for the conditions (standard, staged and control)

METHODS

The intervention was evaluated using a quasi-experimental design with 5 measurement time-points over a period of 2 years.

Study design

Results from randomised control trials are regarded as the gold standard in the hierarchy of research designs. However, research away from laboratories with real people in societies and social structures provides many challenges for evaluating the effectiveness of an intervention. Research in organizations makes it virtually impossible to randomly allocate employees to different conditions, and a pragmatic approach to this intervention was adopted. WWW investigated whether an occupational physical activity intervention can be tailored to target health information according to an individual's readiness to change (staged intervention condition), and if this approach would be more effective than providing standard information (standard intervention condition), or no information at all (control condition). The 10 worksites were allocated to 1 of 3 conditions: staged intervention, standard intervention or control group (to ensure that no cross-contamination of material was possible between employees).

In the staged (tailored) condition the health information was tailored according to recipients' readiness for change. Individuals thinking about increasing their levels of physical activity (contemplation/preparation) were provided with leaflets describing the benefits of physical activity and offering practical tips to increase daily levels of walking. Those not thinking about increasing their physical activity levels (precontemplation) were targeted with leaflets containing awareness raising information about the risks of inactivity. In the standard condition, participants received leaflets which offered generic physical activity advice already available via health promotion organisations. Those in the control group received no intervention material.

For the 2 intervention conditions (staged intervention and standard intervention), in addition to the leaflets administered at baseline and at 6 months, employees were encouraged to increase physical activity levels during the 12-month period via a series of themes introduced every few months. These themes included step count competitions, stair climbing, active commuting (themes were communicated to employees via emailed posters) and a new innovation entitled Walking Lunch. This involves placing a large map (1 metre diameter pinboard) on the wall in a communal area of a work site (e.g. reception area, break room, etc.). The map has a radius of 1.5km and displays the surrounding areas of each work site, which is located in the centre of the map. Employees were encouraged to explore areas on the map to find cafés and restaurants, parks and picnic spots, commuting and walking locations, and places of interest, and record this information on multi-coloured tags pinned to the corresponding location on the map. The activity themes and Walking Lunch were received by both staged and standard conditions, the only factor that differed between the 2 conditions was the provision of information at baseline and at 6 months being either staged/tailored information or standard/generic advice.

Sample

Ten work sites across the UK participated in the study. A large private sector telecommunications organization selected 8 of its work sites, while a medium size public sector local authority involved both of its work sites. Employees at each site were emailed an invitation to participate prior to the recruitment visit (baseline measurement), which contained study information and participant requirements. Posters were also placed around work sites and announcements were made via newsletters. Employees were encouraged to participate in the study with the offer of a free pedometer and feedback from an independent health assessment. Participants completed a questionnaire and undertook physiological and psychological measurements, repeated across: baseline, 6 months, 12 months, 18 months and 24 months.

Sample size

A power calculation indicated that 180 participants would be required in each of the 3 conditions: tailored, standard and control, giving a total sample of 540. The sample size was based on a power calculation for a 3 condition, case-controlled study. This sample size would be necessary to detect a reduction in BMI of 0.3 kg/m². This level of difference was chosen based on a study by Haines et al. (2007) who evaluated the effects of a worksite programme promoting walking. Anticipating a high attrition rate, the study recruited more than double the sample size indicated by the power calculation achieving a sample size of 1120.

Questionnaire measures

A questionnaire was used to collect self-report data on psychological outcomes, physical activity levels and sitting time. The questionnaire was available for participants in paper format or online. The beginning of the questionnaire included a foreword, which on the baseline health screening assessment provided participants with an introduction to the aims of the research. On subsequent revisits, the introduction was modified to notify employees that the questionnaire was only to be completed by those who had already been recruited and were participating in the research. This notification ensured any employees who were not taking part in the research, but may have been forwarded the web-link by a colleague, would not complete the questionnaire.

Participants were asked to record their name and email address at the beginning of the questionnaire. These details were important because the intervention materials, notifications of future health screenings and the Internet web-link to complete questionnaires for future revisits were sent via email. On the questionnaire for the revisits, participants were requested to ensure they recorded the same email address that was used to contact them by the researcher. The email address was also used to match responses for each participant

with their results from previous readings. Each email address was allocated to an identification number, which ensured the results remained confidential as only the researchers had access to these details. The questionnaire comprised ten sections:

- demographic characteristics (age, gender, Office for National Statistics [ONS] job categories, and weekly and monthly hours worked)
- evaluation of participants' readiness to change their physical activity levels
- Domain Specific Sitting Time Questionnaire (Miller and Brown, 2004; Marshall, Miller, Burton and Brown, 2010)
- International Physical Activity Questionnaire (IPAQ) (Craig et al, 2003)
- Occupational Physical Activity Questionnaire (OPAQ) (Reis et al., 2005)
- Work Ability Index (WAI) (Tuomi et al., 1998)
- General Health Questionnaire 12 (GHQ) (Goldberg & Williams, 1988)
- Organizational Commitment (OC) scale (Cook & Wall, 1980)
- Job Motivation (JM) scale (Warr et al., 1979)
- Job satisfaction (JS) and Intention to Quit (ITQ) scales from the Michigan Organizational Assessment Questionnaire (Cammann et al., 1979).

The methods used to score the above scales are described in the companion paper Kazi et al. (submitted).

Stage of change was assessed via a series of yes/no questions:

- 1) Are you planning to increase the amount of physical activity/exercise you do?
 - a. If yes, are you planning to increase the amount of physical activity/exercise you do within the next 6 months?
 - b. If yes, are you planning to increase the amount of physical activity/exercise you do within the next month?

2) Have you recently increased your levels of physical activity/exercise?

- a. If yes, did you make this change (within the last 6 months/more than 6 months ago)

Those responding 'yes' to question 1 and 'yes' to question 1a and 1b were categorised as in the preparation stage. Those responding 'yes' to 1 and 'yes' to 1a but 'no' to 1b were classed as in contemplation. Those responding 'no' to 1 and 'no' to 2 were classed as in pre-contemplation, providing their reported levels of activity were not meeting current activity guidelines. Those responding 'no' to 1 and 'no' to 2 were classed as in maintenance if their reported levels of activity were meeting activity guidelines. Those responding 'no' to 1 and 'yes' to 2 were classed as in action if they made the change less than 6 months ago or maintenance if they had made the change more than 6 months ago.

Self-reported physical activity levels were recorded using the IPAQ short version. Research has shown self-reported physical activity using the IPAQ is comparable to results using objective criterion instruments such as accelerometers (Bauman et al., 2009). The IPAQ has also been used as an outcome measure in previous intervention research (Ferreira et al., 2005). The scoring protocol for the IPAQ short was followed, which was downloaded from the IPAQ website (sites.google.com/site/theipaq).

Physical activity at work was measured using a modified version of the Occupational Physical Activity Questionnaire (OPAQ). The OPAQ is a 7-item measure that identifies the average time per week spent in three occupational activity categories: (a) sitting or standing; (b) walking; and (c) heavy labour. For each category, participants were asked if they performed any of these activities and if they did, to identify the number of hours they performed each activity during a usual working week. For the purposes of this questionnaire, the question that assessed sitting or standing activities at work was edited to read standing activities at work. Sitting time at work was omitted because data on sitting time at work was

collected by the Domain Specific Sitting Time Questionnaire. Participants were also asked to indicate the distance they travelled to work and their usual method of travel to work.

Physiological measurements

An accurate measure of height (in centimetres) was required at each measurement time-point in order to calculate BMI. Height was measured (without shoes) using the Leicester Height Measure, which is a portable plastic stadiometer consisting of a footplate, four-piece vertical ruler and a movable head.

Body weight, BMI and %body fat were measured using a Tanita Body Composition Analyser (Tanita UK Ltd, Model: BC-418 MA, Middlesex, UK) that measures body fat using 8-point bio-impedance analysis. Percentage body fat measured using the Tanita BC-418 has been shown to correlate highly with the reference measure of dual-energy X-ray absorptiometry (DXA) (Pietrobelli et al., 2004). Waist circumference was assessed using anthropometric tape at the midpoint between the upper edge of the iliac crest and the inferior border of the last palpable rib.

Resting blood pressure and heart rate were measured using the validated Omron Intellisense M7 Upper Arm monitor. Two readings were taken after a period of quiet sitting; each reading was separated by a minimum of 30 seconds and the mean of the two readings used in the analyses. If the readings were significantly different, a third reading was taken to collect a more accurate average. If any abnormal readings were identified (e.g. high blood pressure), participants were provided with a referral letter that requested them to visit their GP for further consultation. If this was identified at the baseline health screening, participants were asked to confirm their GP agreed to their participation in this research, and GPs were asked to send a signed confirmation letter to the researchers.

Data handling and analyses

Access scripting was used to match responses using the participant identification number and all data were imported into SPSS Statistics (v22.0) for analyses. To investigate statistically significant differences between all 5 measurement time-points for each intervention condition, linear mixed-model analyses were used to explore significant changes over time.

RESULTS

There was a high level of attrition between baseline and mid-intervention measurements which then stabilised for the return health assessment visits. Return rates, calculated as a percentage from the baseline number recruited were as follows: 33.2% at mid-intervention; 22% end of intervention; 19.1% at 18 months; 14.4% at 24 months. There were several issues (e.g. organizational restructuring, site changes, security issues, etc.) that affected the practical delivery of the intervention in some worksites. These issues had the potential to negatively impact on employees' interest to continue in the research and therefore may have had an impact on return rates.

Table 1 shows the demographic characteristics of the participants, including results from t-tests used to identify gender differences, and one-way analyses of variances (ANOVA) used to explore differences between participants in the 3 conditions. Independent t-tests demonstrated that average height, weight, weekly hours worked, monthly hours worked, and total metabolic equivalent of task (MET) minutes per week were significantly greater in males in comparison to females.

Table 1: Demographic means and SDs[†] for the total sample at baseline, plus gender, ethnicity, and marital status based on allocation to each intervention condition.

Means \pm SD								
	Total sample	Male	Female	P value t-tests	Standard	Staged	Control	P value ANOVA
Age (years)	42.2 \pm 10.3	42.3 \pm 10.4	41.63 \pm 10.4	0.262	43.7 \pm 10.1	42.1 \pm 10.5	39.6 \pm 9.9	0.001
Height (cm)	170.7 \pm 9.8	177.3 \pm 7.2	163.2 \pm 6.6	0.001	172.3 \pm 9.9	169.4 \pm 9.9	170.9 \pm 9.6	0.001
Weight (kg)	78.3 \pm 16.3	84.7 \pm 14.6	70.9 \pm 15.1	0.001	80.8 \pm 16.1	75.3 \pm 16.3	79.8 \pm 15.4	0.001
Weekly hours	36.4 \pm 5.6	37.6 \pm 4.6	35.0 \pm 6.3	0.001	36.0 \pm 5.4	36.7 \pm 5.7	36.7 \pm 5.4	0.078
Monthly hours	147.6 \pm 39.5	154.7 \pm 36.8	139.1 \pm 40.8	0.001	146.9 \pm 36.3	148.2 \pm 41.7	147.3 \pm 40.9	0.935
Total METs p/wk	1826 \pm 1745	2058 \pm 1868	1557 \pm 1550	0.001	1823 \pm 1817	1865 \pm 1653	1749 \pm 1801	0.720
		Standard		Staged		Control		
		n	%	n	%	n	%	
Gender	Male	267	61.9	211	44.8	123	56.4	
	Female	164	38.1	260	55.2	95	43.6	
Marital status	Single	79	18.4	121	25.7	47	21.8	
	Married	264	61.5	233	49.6	111	51.4	
	Civil Partnership	2	0.5	4	0.9	2	0.9	
	Cohabiting	57	13.3	72	15.3	42	19.4	
	Separated	9	2.1	13	2.8	3	1.4	
	Divorced	18	4.2	25	5.3	10	4.6	
	Widowed	0	0.0	2	0.4	2	0.5	
Ethnicity	White ¹	398	93.3	389	83.1	178	82.4	
	Asian ²	20	4.7	50	10.8	31	14.4	
	Black ³	7	1.5	20	4.2	4	1.8	
	Mixed ⁰	2	0.5	9	1.9	3	1.4	

[†]Table includes significance values of the t-tests and ANOVA assessing gender and intervention group differences

¹White (British, Irish, other) ²Asian or Asian British (Indian, Pakistani, Bangladeshi, Chinese, other)

³Black or Black British (Caribbean, African, other) ⁰Mixed (White and Asian, White and Black, other mixed)

Table 2 summarises the health outcomes for each intervention group at each measurement time-point during and post intervention

Accepted Manuscript

Table 2: Means±SDs for each intervention group (standard, staged, control) at each time-point during and post intervention.

		Baseline			Mid-intervention			End of intervention			6m Follow up			12m Follow up		
		Stan dard	Sta ged	Con trol	Stan dard	Sta ged	Con trol	Stan dard	Sta ged	Con trol	Stan dard	Sta ged	Con trol	Stan dard	Sta ged	Con trol
Physiological	BMI	27.2 ± 4.6	26.2 ± 5	27.2 ± 4.8	26.9 ± 4.1	25.5 ± 4.6	26.6 ± 4.2	26.0 ± 4.0	25.6 ± 4.5	26.6 ± 3.8	26.1 ± 3.6	25.2 ± 4.5	26.3 ± 3.2	26.1 ± 3.6	24.7 ± 3.8	25.5 ± 3.6
	Fat %	28.4 ± 8.9	29.1 ± 9.1	29.4 ± 9.4	26.4 ± 8.2	26.9 ± 8.1	26.7 ± 9.1	25.7 ± 8.6	26.8 ± 8.3	25.1 ± 8.2	25.4 ± 8.3	27 ± 8.0	27.7 ± 9.4	24.0 ± 8.0	25.6 ± 7.5	27.7 ± 9.5
	WC	92.9 ± 13.2	88.1 ± 13.7	91.9 ± 12.5	93.4 ± 12.0	88.0 ± 14.0	91.2 ± 13.3	90.7 ± 11.7	87.7 ± 14.0	91.4 ± 12.2	91.3 ± 10.4	86.9 ± 14.2	89.2 ± 9.5	91.5 ± 11.2	85.3 ± 12.7	87.0 ± 12.9
	SBP	131.6 ± 16.4	128. ± 6	129. ± 15.1	132.0 ± 15.2	124. ± 16.9	128. ± 13.4	133.1 ± 14.6	130. ± 17.2	132. ± 12.4	133.4 ± 14.9	130. ± 16.3	132. ± 13.9	130.1 ± 13.0	126. ± 16.5	132. ± 16.8
	DBP	79.2 ± 10.4	77.8 ± 10.1	77.4 ± 10.2	81.1 ± 11.2	75.6 ± 10.6	76.4 ± 9.7	78.2 ± 9.4	77.6 ± 10.6	76.5 ± 10.3	77.9 ± 9.8	76.9 ± 9.2	79.9 ± 11.5	75.4 ± 9.3	75.2 ± 9.7	79 ± 10.9
	HR	66.6 ± 11.0	67.0 ± 10.8	69.3 ± 12.6	64.7 ± 11.0	64.8 ± 11.8	64.8 ± 10.5	64.8 ± 11.7	67.1 ± 12.1	63.0 ± 11.1	65.8 ± 12.9	66.0 ± 14.3	64.0 ± 13.6	63.0 ± 11.0	66.2 ± 12.9	64.5 ± 13.6
Psychological	WAI	42.4 ± 4.1	42.3 ± 4.7	41.7 ± 4.9	32.4 ± 10.8	31.1 ± 12.1	30.6 ± 12.3	42.7 ± 4.5	42.8 ± 3.8	41.7 ± 5.4	36.3 ± 15.9	32.9 ± 18.6	17.0 ± 21.5	43.3 ± 3.8	41.9 ± 4.4	42.9 ± 2.0
	OC	10.7 ± 4.9	11.1 ± 5.1	11.6 ± 5.2	9.7 ± 5.6	10.2 ± 5.8	10.3 ± 6.3	9.9 ± 5.3	11.8 ± 5.5	11.2 ± 5.6	8.6 ± 5.8	7.9 ± 6.0	6.1 ± 7.2	9.9 ± 3.6	11.5 ± 5.2	9.1 ± 4.3
	JM	5.4 ± 1.2	5.3 ± 1.3	5.0 ± 1.4	5.3 ± 1.2	5.3 ± 1.3	5.2 ± 1.3	5.4 ± 1.3	5.3 ± 1.4	5.3 ± 1.3	4.5 ± 2.2	± 2.5	± 3.1	5.3 ± 1.3	± 1.4	± 1.6
	ITQ	46.4 ± 7.8	46.6 ± 8.1	43.6 ± 8.8	41.0 ± 4.4	39.6 ± 5.3	39.9 ± 5.3	46.4 ± 7.3	45.7 ± 8.1	45.0 ± 8.4	38.8 ± 17.7	36.7 ± 20.9	24.4 ± 24.6	46.1 ± 6.9	46.0 ± 9.1	43.1 ± 12.1
	GHQ	34.8 ± 3.7	35.1 ± 4.0	34.0 ± 4.1	31.3 ± 10.8	30.4 ± 11.6	30.1 ± 12.2	34.9 ± 4.0	± 4.0	± 5.4	29.5 ± 12.7	27.0 ± 14.9	17.6 ± 17.5	34.1 ± 4.4	34.0 ± 5.0	32.6 ± 5.4
	JS	2.7 ± 1.5	2.9 ± 1.6	3.4 ± 1.9	2.5 ± 1.6	2.4 ± 1.9	2.6 ± 1.9	2.4 ± 1.3	2.9 ± 1.6	2.9 ± 1.5	2.3 ± 1.7	1.8 ± 1.6	1.1 ± 1.3	2.6 ± 1.4	2.8 ± 1.4	3.3 ± 1.9
MET-mins	Walki ng	752.0 ± 903.4	888. ± 825. 9	802. ± 868. 4	1029. ± 1219. 1	801. ± 927. 9	748. ± 900. 6	1151. ± 1305. 7	137 ± 166 9	818. ± 908. 6	1334. ± 3343. 1	167 ± 477 7.3	183 ± 597 8.0	1086. ± 1321. 4	149 ± 192 7.2	462. ± 517. 2
	Mode rate PA	338.8 ± 676.1	290. ± 620. 2	221. ± 470. 4	468.1 ± 828.5	302. ± 679. 7	551. ± 113 5.4	437.1 ± 906.1	439. ± 101 3.2	365. ± 787. 2	456.2 ± 860.4	616. ± 368 7.4	535. ± 139 7.1	767.8 ± 1427. 5	543. ± 126 4.1	114 ± 164 6.7
	Vigor ous PA	733.4 ± 1164. 1	685. ± 105 4.3	725. ± 116 0.8	870.0 ± 1393. 5	834. ± 111 7.9	875. ± 176 7.4	802.3 ± 1419. 8	789. ± 110 1.6	120 ± 172 7.6	860.3 ± 2320. 3	105 ± 177 6.6	998. ± 232 5.5	814.9 ± 1180. 2	886. ± 157 5.3	269 ± 352 0.0
Workday sitting time	Trans port	56.1 ± 57.8	59.8 ± 43.9	46.4 ± 52.9	42.2 ± 38.9	56.1 ± 43.9	41.1 ± 38.4	47.6 ± 41.6	61.1 ± 42.2	43.6 ± 36.1	42.9 ± 36.2	49.9 ± 47.1	34.8 ± 46.3	58.9 ± 80.9	62.0 ± 41.0	67.5 ± 56.8
	Work	372.4 ± 125.9	373. ± 125. 2	385. ± 107. 0	345.2 ± 151.3	343. ± 149. 2	330. ± 89.2	381.6 ± 110.7	395. ± 89.5	376. ± 115. 8	337.7 ± 165.1	301. ± 176. 9	168. ± 184. 7	387.5 ± 80.9	375. ± 79.9	352. ± 74.8
	TV	94.7 ± 64.7	89.9 ± 70.0	103. ± 69.4	89.3 ± 61.8	79.8 ± 62.3	89.2 ± 62.3	98.1 ± 68.9	84.5 ± 61.9	107. ± 67.3	81.7 ± 67.4	77.4 ± 67.5	38.4 ± 62.2	122.6 ± 82.7	92.7 ± 60.7	120. ± 90.7
	Hom	51.7	58.9	65.6	50.2	56.7	54.2	60.9	64.6	52.4	45.1	44.7	28.4	60.0	43.3	62.5

	e PC	± 74.6	± 101. 0	± 93.1	± 67.3	± 102. 4	± 86.7	± 84.2	± 102. 5	± 66.3	± 74.3	± 76.2	± 73.6	± 75.9	± 60.0	± 85.2
	Leisure	43.8 ± 56.8	46.5 ± 65.8	39.4 ± 51.9	35.8 ± 53.4	38.0 ± 51.2	37.6 ± 51.7	42.5 ± 55.5	51.9 ± 58.0	42.6 ± 60.6	39.9 ± 51.0	32.7 ± 48.3	27.4 ± 48.8	39.9 ± 48.5	51.3 ± 55.8	30.0 ± 45.4
	Total	603.0 ± 152.2	599. 7 ± 162. 3	615. 9 ± 142. 9	615.1 ± 115.5	613. 4 ± 145. 6	633. 8 ± 135. 8	607.4 ± 144.2	616. 6 ± 133. 9	615. 9 ± 184. 5	608.3 ± 124.0	625. 4 ± 122. 3	531. 7 ± 199. 4	648.0 ± 134.7	604. 6 ± 138. 2	632. 5 ± 145. 5

BMI: Body Mass Index; Fat %: Fat Percentage; WC: Waist Circumference; SBP: Systolic Blood Pressure; DBP: Diastolic Blood Pressure; HR: Resting Heart Rate; WAI: Work Ability Index; OC: Organizational Commitment; JM: Job Motivation; ITQ: Intention to Quit; GHQ: General Health Questionnaire; JS: Job Satisfaction; PA: Physical Activity.

Accepted Manuscript

Low numbers of returning control participants at the 6 month post intervention point resulted in a fluctuation in the mean WAI score. The mean WAI was reduced by several respondents with particularly low scores at this measurement point.

Table 3 displays the results from the linear mixed-model analyses assessing the effects of the standard and staged intervention conditions when compared to the control group (intercept).

Accepted Manuscript

Table 3: Linear mixed model test of fixed effects for predicting impact of standard/staged intervention condition. [†]

		Intercept				
		Control	Standard	Sig.	Staged	Sig.
Physical	BMI	27.15	-0.06	0.882	-1.05	0.007*
	Fat %	29.28	-1.15	0.127	-0.31	0.672
	WC	91.73	0.99	0.362	-3.88	0.001**
	SBP	129.08	1.97	0.115	-1.52	0.220
	DBP	77.14	1.50	0.064	-0.04	0.961
	HR	68.64	-2.10	0.021*	-1.78	0.047*
Psychological	WAI	41.45	0.82	0.029*	0.77	0.039*
	OC	42.87	2.52	0.001**	2.64	0.001**
	JM	33.70	0.96	0.003*	1.20	0.001**
	ITQ	3.16	-0.53	0.001**	-0.32	0.010*
	GHQ	11.18	-0.80	0.033*	-0.20	0.590
	JS	4.99	0.34	0.001*	0.28	0.007*
MET-mins	Walking	797.09	68.64	0.780	67.95	0.091
	Moderate PA	273.05	88.99	0.065	27.27	0.568
	Vigorous PA	784.61	-38.39	0.667	-78.88	0.371
Workday sitting time	Transport	44.57	6.57	0.29*	13.93	0.001**
	Work	357.72	7.33	0.376	5.96	0.476
	TV	97.22	-2.85	0.495	-10.36	0.014*
	Home PC	58.89	-6.73	0.208	-2.24	0.678
	Leisure	38.47	2.92	0.406	6.14	0.084
	Total	616.70	-7.23	0.456	-11.09	0.260

[†] Reported as estimates of mean parameters (standard error).

*P<0.05, **p<0.001

The staged intervention group showed a significant reduction in BMI over the duration of the intervention period, including post intervention in comparison to the standard intervention group and control group (intercept). The results show that participants provided with tailored health information were more likely to have reduced BMI by -1.05 kg/m^2 over the course of the measurement period.

There were no significant differences in Fat% for either of the intervention conditions when compared to the control group. This demonstrates that even though a reduction in BMI was observed, this was most likely due to an overall weight reduction rather than specifically fat reduction. In contrast, waist circumference was significantly lower for the staged intervention group in comparison to the standard intervention group, which may indicate reductions in visceral fat.

Table 3 also shows there were no significant differences in blood pressure outcomes (systolic or diastolic) for the duration of the intervention period in either of the intervention groups. However, for both intervention groups, there were significant improvements (reductions) in resting heart rate.

In terms of psychological outcomes, significant differences in both intervention groups (standard and staged) were seen in comparison to the control group. The results show that employees receiving either standard or tailored intervention material demonstrated significantly higher self-reported work ability, organizational commitment, job motivation, job satisfaction, and a reduction in intention to quit the organization.

While both the staged and standard interventions reduced total sitting times compared to controls, the differences failed to reach significance. The results showed a significant increase in sitting time for both standard and staged groups for the domain of transport in

comparison to the control. However, the staged intervention group showed a significant reduction in sitting time while watching TV compared to the control group.

Table 4 shows the percentage of participants in each stage of change by intervention condition and at each stage of the intervention and follow up.

Table 4: Stage of change classifications by intervention condition at each stage of the intervention and follow up (%).

Intervention Condition	Stage of Change	Baseline	Mid-intervention	End of intervention	6m Follow up	12m Follow up
Standard	Precontemplation	20.9	21.6	28.4	20.2	30.5
	Contemplation	14.5	11.2	15.9	13.5	11.9
	Preparation	54.0	46.2	34.1	49.4	33.9
	Action	9.7	4.2	9.1	9.0	20.3
	Maintenance	0.9	16.8	12.5	7.9	3.4
Staged	Precontemplation	16.7	19.4	20.9	25.0	26.9
	Contemplation	16.4	7.4	20.8	12.4	9.7
	Preparation	55.9	42.6	34.7	39.1	36.5
	Action	8.1	13.0	4.2	18.8	19.2
	Maintenance	2.9	17.6	19.4	4.7	7.7
Control	Precontemplation	14.5	21.1	23.5	23.1	25.0
	Contemplation	13.6	8.5	13.7	15.3	12.5
	Preparation	56.1	42.3	51.0	23.1	50.0
	Action	13.5	4.2	2.0	30.8	12.5
	Maintenance	2.3	23.9	9.8	7.7	0.0

For participants in the standard intervention condition, a chi-squared goodness of fit test indicated significant differences in the proportion distribution of participants for each stage at mid-intervention and end of intervention measurements compared to baseline. At 6 months there were fewer workers in contemplation and action and more participants in the maintenance stage ($\chi^2=70.68$, $p<0.001$). By 12 months there were fewer employees in preparation and more in the maintenance stage in comparison to baseline ($\chi^2=30.20$, $p<0.001$).

For participants in the staged intervention condition, a chi-squared goodness of fit test indicated that at 6 months there were fewer workers in contemplation and more in maintenance ($\chi^2=22.31$, $p<0.001$). At 12 months there were fewer workers in preparation and more in maintenance in comparison to baseline ($\chi^2=30.20$, $p<0.001$).

For participants in the control condition, a chi-squared goodness of fit test indicated that at 6 months there were fewer participants in preparation and more in maintenance ($\chi^2=40.30$, $p<0.001$). However, these differences were not maintained by the end of intervention, as there was no significant difference in the proportion of employees for each stage of change compared to the baseline assessment ($\chi^2=5.16$, $p<0.271$).

DISCUSSION

This longitudinal study aimed to evaluate the effectiveness of a tailored workplace physical activity intervention compared to a standard condition and control condition implemented in 10 different worksites across the UK. The results showed that the tailored intervention group demonstrated significant reductions in BMI over the duration of the 12 month intervention period and the 12 month follow-up period in comparison to the standard intervention and control group. Additionally, waist circumference was significantly lower for the staged intervention group in comparison to the standard intervention and control group, which is indicative of reductions in visceral fat. The findings provide new evidence that tailored interventions are more effective in improving health outcomes in the workplace. These results support the calls for adopting ergonomics interventions which align with workers' stage of change (Haslam and Haslam, 2001; Haslam, 2002),

There were no significant differences in blood pressure outcomes for the duration of the intervention period in either of the intervention groups. However, for both intervention groups, there were significant improvements (reductions) in resting heart rate. Employees who received either standard or tailored intervention material demonstrated significantly higher self-reported work ability, organizational commitment, job motivation, job satisfaction, and a reduction in intention to quit the organization. These results suggest that physical activity interventions have a positive impact on employees' job attitudes and psychological wellbeing.

Both staged and standard interventions reduced total sitting times, but the differences failed to reach significance. Both conditions showed a significant increase in sitting times during transport (a domain where employees may have limited control) whereas the staged intervention demonstrated a significant reduction in sitting times while watching TV (where they would have control).

A strength of this study is that it was longitudinal, and the intervention was evaluated over a 2 year period incorporating a wide range of objective physical measurements as well as self-reported health and psychological outcomes. The companion paper Kazi et al. (submitted) reporting the findings of the baseline phase of this study recommends targeting future interventions to according to gender, job role, sector and geographical location. This was not possible in this instance as this research was the longitudinal phase of the study and was constrained by the sample recruited at baseline. As work sites were allocated to intervention conditions it was not possible to match the samples across the conditions. The tailored condition comprised more female participants (55%) compared to control and standard conditions (43% and 38% respectively) and this may have contributed to the outcomes. Another limitation of the study was the initial high level of attrition from baseline to mid-intervention. This was due to a number of organisational changes that were outside the control of the study. Such factors are inherent in organisational research. The attrition levels stabilised for the additional visits which enabled sufficient samples in all conditions to conduct the long-term evaluation.

This paper describes the development and evaluation of a new intervention, designed to increase physical activity and reduce sedentary behaviour in the workplace. The intervention involves tailoring health information to the individual employee. This approach assumes that when attempting to change behaviour, one size does not fit all, and success is greater when interventions align with peoples' beliefs. This longitudinal study, conducted across 10 worksites compared tailored, standard and control conditions and showed that while both tailored and standard interventions reduced resting heart rate and improved psychological wellbeing, the tailored approach was more effective in reducing BMI and waist circumference compared to standard information and control conditions.

Sedentary work is a major public health concern with high levels of sitting associated with an increased risk of premature mortality and a wide range of chronic health conditions. This

study has shown that interventions tailored to employees' stage of change are more effective in reducing BMI and waist circumference than standard approaches and this is an important finding. Kearns et al. (2014) investigated the burden of chronic disease associated with overweight and obesity in the adult population and estimated that a 1 unit reduction in BMI at a population level would lead to substantial gains in terms of reduced prevalence of chronic diseases. The present study demonstrated a reduction in BMI of 1.05 kg/m² which suggests that tailored workplace physical activity interventions have the potential to reduce the burden of chronic diseases.

We would argue that sitting at work should be considered alongside other risk exposures in the workplace. Where employees are spending protracted periods of sitting at work, risk assessments should be undertaken to determine what measures may be put in place to ameliorate this important health risk. Detailed risk assessments may allow occupational health initiatives to better target physical activity interventions to job roles and sectors where sedentary behaviour is more prevalent.

The workplace is an ideal arena for targeting health information and intervening to improve health and wellbeing. The baseline results of this longitudinal study reported in the companion paper, Kazi et al. (submitted) have highlighted important gender, job role, sector and geographical differences in sedentary behaviour in the workplace. This information may be used to inform future workplace health initiatives enabling interventions to be more specifically targeted and more effective in improving health outcomes. The results of this present study indicate that tailoring information to employees' stage of change is more effective than standard approaches in improving health in the workplace. These results provide important insights for future workplace physical activity interventions and offer considerable scope to counteract increasing levels of obesity and improve the health and wellbeing of the sedentary workforce.

References

- Andersen, S., C. Keller. 2002. "Examination of the Transtheoretical Model in Current Smokers." *Western Journal of Nursing Research*, 24 (3): 282–294.
- Bauman, A., B. E. Ainsworth, F. Bull, C. L. Craig, M. Hagströmer, J. F. Sallis, M. Pratt and M. Sjöström. 2009. "Progress and pitfalls in the use of the International Physical Activity Questionnaire (IPAQ) for adult physical activity surveillance." *Journal of Physical Activity and Health*, 6(Suppl. 1): S5–S8.
- Biswas, A., P. I. Oh, G. E. Faulkner, R. R. Bajaj, M. A. Silver, M. S. Mitchell, and D. A. Alter. 2015. "Sedentary Time and its Association with Risk for Disease Incidence, Mortality, and Hospitalization in Adults: A Systematic Review and Meta-Analysis." *Annals of Internal Medicine* 162 (2): 123-132.
- Cammann, C., M. Fichman, D. Jenkins and J. Klesh. 1979. The Michigan Organizational Assessment Questionnaire. University of Michigan: Unpublished manuscript.
- Chau, J. Y., A. C. Grunseit, T. Chey, E. Stamatakis, W.J. Brown, C. E. Matthews, A. E. Bauman, and H. P. van der Ploeg. 2013. "Daily sitting time and all-cause mortality: a meta-analysis." *PLoS ONE* 8 (11): e80000.
- Cook, J. and T. Wall. 1980. "New work attitude measures of trust, organizational commitment and personal need non-fulfilment." *Journal of Occupational Psychology* 53 (1): 39–52.
- Craig, C. L., A. L. Marshall, M. Sjöström M, A. E. Bauman, M. L. Booth, A. E. Ainsworth, M. Pratt, U. Ekelund, A. Yngve, J. F. Sallis, P. Oja. 2003. "International physical activity

questionnaire: 12-country reliability and validity.” *Medicine and Science in Sports and Exercise* 35 (8):1381–1395.

Ferreira, M., S. Matsudo, V. Matsudo and G. Braggion. 2005. “Effects of an intervention program of physical activity and nutrition orientation on the physical activity level of physically active women aged 50 to 72 years old.” *Revista Brasileira de Medicina do Esporte* 11: 166–169.

Gierach, G. L., S. C. Chang, L. A. Brinton, J. V. Lacey Jr, A. R. Hollenbeck, A. Schatzkin, and M. F. Leitzmann. 2009. “Physical activity, sedentary behavior, and endometrial cancer risk in the NIH-AARP Diet and Health Study.” *International Journal of Cancer* 124 (9): 2139–2147.

Goldberg, D. P. and P. Williams. 1988. *A user’s guide to the General Health questionnaire*. Windsor, UK: NFER-Nelson.

Haines, D. J., L. Davis, R. Rancour, M. Robinson, T. Neel-Wilson and S. Wagner. 2007. “A Pilot Intervention to Promote Walking and Wellness and to Improve the Health of College Faculty and Staff.” *Journal of American College Health* 55 (4): 219-225.

Hamilton, M. T., D. G. Hamilton, and T. W. Zderic. 2007. “Role of low energy expenditure and sitting in obesity, metabolic syndrome, type 2 diabetes, and cardiovascular disease.” *Diabetes* 56 (11): 2655–2667.

Haslam, C. 2000. “A targeted approach to reducing maternal smoking.” *British Journal of General Practice* 50: 661-663.

Haslam, C. and E. S. Draper. 2000. “Stage of change is associated with assessment of the health risks of maternal smoking among pregnant women.” *Social Science and Medicine* 51:

1189-1196.

Haslam, C. and R. Haslam. 2001. "A stage specific approach to improving occupational health and safety." In: Hanson, M.A. (Ed.), *Contemporary Ergonomics 2001*. Taylor & Francis, London, pp.171– 176.

Haslam, C. and W. Lawrence. 2004. "Health related behavior and beliefs of pregnant smokers." *Health Psychology* 23 (1): 486-491.

Haslam, R. A. "Targeting ergonomics interventions – learning from health promotion." *Applied Ergonomics* 33: 241-249.

Hu, G., Q. Qiao, K. Silventoinen, J. G. Eriksson, P. Jousilahti, J. Lindström, J. T. T. Valle, A. Nissinen and J. Tuomilehto, J. 2003. "Occupational, commuting, and leisure-time physical activity in relation to risk for Type 2 diabetes in middle-aged Finnish men and women." *Diabetologia* 46 (3): 322–329.

International Physical Activity Questionnaire. sites.google.com/site/theipaq. Accessed 24 November 2015.

Katzmarzyk, P. T., T. S. Church, C. L. Craig and C. Bouchard. 2009. "Sitting time and mortality from all causes, cardiovascular disease, and cancer." *Medicine & Science in Sports & Exercise* 41 (5): 998-1005.

Kazi, A. 2013. Promoting physical activity in the workplace: a stage of change approach. PhD thesis, Loughborough University.

Kazi, A., M. Duncan, S. Clemes and C. Haslam. 2014. "A survey of sitting time among UK

employees." *Occupational Medicine* 64 (7) 497-502.

Kazi, A. C. Haslam, M. Duncan, S. Clemes, R. Twumasi. "Sedentary behaviour and health at work: an investigation of industrial sector, job role, gender and geographical differences."

Submitted to *Ergonomics*.

Kearns, K., A. Dee, A. P. Fitzgerald, E. Doherty and I. J. Perry. 2014. "Chronic disease burden associated with overweight and obesity in Ireland: the effects of a small BMI reduction at population level." *BMC Public Health* 14: 143.

Kirk, A., N. Mutrie, P. MacIntyre and M. Fisher, M. 2003. "Increasing physical activity in people with type 2 diabetes." *Diabetes Care* 26 (4): 1186–1192.

Lynch, B. M. 2010. "Sedentary behavior and cancer: a systematic review of the literature and proposed biological mechanisms." *Cancer Epidemiology Biomarkers & Prevention* 19 (11): 2691–2709.

Mansoubi, M., N. Pearson, S. J. H. Biddle, and S. A. Clemes. 2014. "The relationship between sedentary behaviour and physical activity in adults: A systematic review." *Preventive Medicine*, 69: 28–35.

Marshall, S. J. and S. J. H. Biddle. 2001. "The transtheoretical model of behaviour change: a meta-analysis of applications to physical activity and exercise." *Annals of Behavioral Medicine*, 23 (4): 229–246.

Marshall, A. L., Y. D. Miller, N. W. Burton, W. N. Brown. 2010. "Measuring total and domain-specific sitting: a study of reliability and validity." *Medicine and Science in Sports and Exercise* 42 (6): 1094–1102.

Miller, R. and W. Brown W. 2004. "Steps and sitting in a working population." *International Journal of Behavioural Medicine* 11 (4): 219–224.

Owen, N., A. Bauman, and W. Brown. 2009. "Too much sitting: a novel and important predictor of chronic disease risk?" *British Journal of Sports Medicine* 43 (2): 81-83.

Pietrobelli, A., F. Rubiano, M. P. St-Onge and S. B. Heymsfield. 2004. "New bioimpedance analysis system: improved phenotyping with whole-body analysis." *European Journal of Clinical Nutrition* 58 (11): 1479-1484.

Povey, R., M. Conner, P. Sparks, R. James and R. Shepherd, R. 1999. "A critical examination of the application of the Transtheoretical Model's stages of change to dietary behaviours." *Health Education Research* 14 (5): 641–651.

Prochaska, J. O. and C. C. DiClemente, C. C. 1982. "Transtheoretical therapy: Toward a more integrative model of change." *Psychotherapy: Theory Research and Practice* 19 (3): 276-288.

Prochaska, J. O. and C. C. DiClemente. 1983. "Stages and processes of self-change of smoking: Toward an integrative model of change." *Journal of Consulting and Clinical Psychology*, 51 (3): 390–395.

Prochaska, J. O., C. C. DiClemente, W. F. Velicer and J. S. Rossi. 1993. "Standardised, individualised, interactive, and personalised self-help programmes for smoking cessation." *Health Psychology* 12 (5): 399-405.

Proper, K. I., A. S. Singh, W. van Mechelen and M. J. M. Chinapaw. 2011. Sedentary

behaviors and health outcomes among adults: a systematic review of prospective studies.”
American Journal of Preventive Medicine 40 (2): 174–182.

Reis, J. P., K. D. Dubose, B. E. Ainsworth, C. A. Macera and M. M. Yore. 2005. “Reliability and Validity of the Occupational Physical Activity Questionnaire.” *Medicine & Science in Sports & Exercise* 37 (12): 2075–2083.

Rothmore, P., P. Aylward, J. Gray and J. Karnon. 2017. “A long-term evaluation of stage of the change approach and compensable injury outcomes – a cluster randomised trial.”
Ergonomics 60 (5): 628-635.

Tuomi, K., J. Ilmarinen, J., A. Jahkola, L. Katajarinne, and A. Tulkki. 1988. Work Ability Index. Work (2nd ed.). Helsinki: Finnish Institute of Occupational Health.

van Uffelen, J. G. Z, J. Wong, J. Y. Chau, H. P. van der Ploeg, I. Riphagen, N. D. Gilson, N. W. Burton, G. N. Healy, A. A. Thorpe, B. K. Clark, P. A. Gardiner, D. W. Dunstan, A. Bauman, N. Owen, and W. J. Brown. 2010. “Occupational sitting and health risks: a systematic review.” *American Journal of Preventive Medicine* 39 (4): 379–388.

Warr, T., P. Cook and K. Wall, K. 1979. “Scales for the measurement of some work attitudes and aspects of psychological well-being.” *Journal of Occupational Psychology* 52 (2): 129–148.

Whysall, Z. J., C. Haslam and R. Haslam. 2005. *A staged approach to reducing musculoskeletal disorders (MSDs)*. Health and Safety Executive. HSE Research Report 379. HSE Books: Sudbury, UK.

Whysall, Z. J., C. Haslam and R. Haslam. 2006. “A staged approach to reducing

occupational ill health.” *Preventive Medicine* 43 (5): 422–428.

Whysall, Z. J., C. Haslam and R. Haslam. 2007. “Developing the stage of change approach for the reduction of work-related musculoskeletal disorders.” *Journal of Health Psychology* 12 (1): 184–197.

Wilmot, E. G., C. L. Edwardson, F. A. Achana, M. J. Davies, T. Gorely, L. J. Gray, K. Khunti, T. Yates, and S. J. Biddle. 2012. “Sedentary time in adults and the association with diabetes, cardiovascular disease and death: systematic review and meta-analysis.” *Diabetologia* 55 (11): 2895–2905.